

Anomalous Negative Magnetoresistance Caused by Non-Markovian Effects

Vadim V. Cheianov¹, A. P. Dmitriev², V. Yu. Kachorovskii²

¹*NORDITA, Blegdamsvej 17, Copenhagen, DK 2100, Denmark*

²*A.F. Ioffe Physical-Technical Institute, 26 Polytechnicheskaya str., Saint Petersburg, 194021, Russia*

(Dated: February 2, 2008)

A theory of recently discovered anomalous low-field magnetoresistance is developed for the system of two-dimensional electrons scattered by hard disks of radius a , randomly distributed with concentration n . For small magnetic fields the magnetoresistance is found to be parabolic and inversely proportional to the gas parameter, $\delta\rho_{xx}/\rho \sim -(\omega_c\tau)^2/na^2$. With increasing field the magnetoresistance becomes linear $\delta\rho_{xx}/\rho \sim -\omega_c\tau$ in a good agreement with the experiment and numerical simulations.

PACS numbers: 05.60.+w, 73.40.-c, 73.43.Qt, 73.50.Jt

It is well known that in the Boltzmann-Drude approach the longitudinal resistivity ρ_{xx} of a degenerate two-dimensional (2D) electron gas does not depend on the transverse magnetic field B . Therefore, the known mechanisms of magnetoresistance (MR) involve either quantum interference effects or classical non-Markovian memory effects, which are not captured in the Boltzmann picture. The MR, arising from quantum effects was discussed in a great number of works (see for review Ref. [1]). The role of classical memory effects was underappreciated for a long time, though several theoretical works pointed out at the importance of such effects for magnetotransport [2, 3, 4]. The interest to the problem of classical MR has sharply increased in recent years, starting with Ref. [5], where it was shown that effects of "classical localization" may lead to the exponential suppression of electron diffusion at large B . This work was followed by a series of works [6, 7, 8, 9, 10, 11, 12, 13, 14], discussing different aspects of classical 2D magnetotransport in strong magnetic fields.

In this paper we focus on a mechanism of low field classical MR specific for systems of strong scatterers. This mechanism is connected to the memory effects due to backscattering events. The corresponding corrections to the conductivity are small in the parameter a/l , where a is the characteristic size of the scatterers and l is the mean free path. Nevertheless, the dependance of these corrections on the magnetic field turns out to be very sharp, resulting in the MR anomaly. The anomaly was discovered in recent numerical simulations [15] where the MR in a system of 2D electrons scattering on randomly distributed hard disks was studied. This system is usually referred to as the Lorenz gas and is the simplest model of the 2D electron gas with strong scatterers. In the following we restrict our considerations to this model. The generalization of our results to other models of strong disorder is straightforward. Magnetotransport in the Lorenz gas is characterized by two dimensionless parameters: $\beta = \omega_c\tau$, and the gas parameter $\beta_0 = a/l = 2na^2$. Here a is the disk radius, n is disks' concentration, ω_c is the cyclotron frequency, $\tau = l/v_F$ is the mean free time and

$l = 1/2na$ is the mean free path. The anomaly was observed in the case $\beta \ll 1$, $\beta_0 \ll 1$. Both the numerical simulations and the qualitative analysis of [15] indicated that at zero temperature, T , the MR can be expressed in terms of a dimensionless function $f(z)$ via

$$\frac{\delta\rho_{xx}}{\rho} = -\beta_0 f\left(\frac{\beta}{\beta_0}\right), \quad (1)$$

where ρ is the resistivity for $B = 0$. Numerical results [15] suggest that $f(z) \sim z$ as $z \rightarrow 0$, yielding $\delta\rho_{xx}/\rho \sim -|\omega_c|\tau$. The latter expression is in a very good agreement with experimental measurements of MR in a random antidot arrays [16]. It is anomalous in two senses. First, it has a non-analytic dependence on the magnetic field. Second, it does not vanish in the limit of vanishing β_0 , which is normally regarded as the expansion parameter for the corrections to the Drude-Boltzmann picture. This intriguing behavior calls for a rigorous analytical theory of the effect, which would establish Eq. 1 and enable one to derive the analytical expression for function f . In this letter we present a theory of the anomaly and give an expression for $f(z)$. We find that for $z \lesssim 1$, $f(z)$ is linear in agreement with numerical experiment, but at very small $z \lesssim 0.05$ crosses over to a quadratic dependence. Thus, for $\beta \rightarrow 0$, Eq. 1 yields $\delta\rho_{xx}/\rho \sim -\beta^2/\beta_0$. The limit $\beta_0 \rightarrow 0$ should be taken with care. While the small β expansion seems to be singular as a function of β_0 , the region of β where this expansion is valid shrinks as $\beta_0 \rightarrow 0$. For $z \rightarrow \infty$, f saturates at some constant value. Therefore, the full variation of $\delta\rho_{xx}/\rho$ is of the order β_0 . In other words, the anomalous MR is strong but it exists in a small region of magnetic fields.

In [15] a mechanism of MR connected with memory effects arising in backscattering events was proposed. It has a close relation to the well known non-analyticity of the virial expansion of transport coefficients [17, 18, 19, 20, 21], which we briefly recall. For $B = 0$ the leading nonanalytic correction to resistivity, $\delta\rho$, is due to the processes of return to a scatterer after a single collision with another scatterer [Fig. 1(a)]. The relative correction, $\delta\rho/\rho$, is proportional to the correspond-




FIG. 1: Backscattering process responsible for leading non-analytic contribution to the resistivity at $B = 0$ (a). For $B \neq 0$, the overlap area, S_B , between two corridors is small at large B (b). For $\phi = 0$, S_B decreases with B (c). For $\phi \neq 0$ and small B the values of $S_B - S_0$ for time reversed trajectories have opposite signs (d,e).

ing backscattering probability, given by the product of $e^{-r/l} d\phi dr/l$ (which is the probability to reach scatterer 2 without collision and scatter in the angle $d\phi$) and the probability p to return without collisions from 2 to 1 (here l is the mean free path). Assuming $p = \exp(-r/l)$ and integrating over intervals $0 < \phi < a/r$, $a < r < \infty$, one obtains [17, 18, 19, 20, 21] $\delta\rho/\rho \sim \beta_0 \ln(1/2\beta_0)$.

In Ref. [15] it was shown that the probability p is actually larger than $\exp(-r/l)$ because the passage of a particle from 1 to 2 ensures the existence of a corridor of width $2a$ free of the centers of the disks. This reduces the scattering probability on the way back, yielding $p(r, \phi) = \exp(-r/l + nS_0(r, \phi))$, where $S_0(r, \phi) = 2ar - r^2|\phi|/2$ is the area of the overlap of the two corridors [Fig. 1(a)]. For example, for $\phi = 0$, we have $S_0 = 2ar$ and $p = 1$. Physically, this means that the particle is unable to scatter, since it travels back along the same path. Taking into account the effect of “empty corridor”, we get

$$\frac{\delta\rho}{\rho} \sim \int_a^\infty \frac{dr}{l} \int_0^{a/r} d\phi e^{-(2r/l) + nS_0} \approx \beta_0 \ln\left(\frac{C}{2\beta_0}\right), \quad (2)$$

where C is a constant of the order of unity. Thus, for $B = 0$ this effect simply changes the constant in the argument of the logarithm.

The key idea suggested in [15] was that for $B \neq 0$ the area of the overlap of the two corridors, S_B , sharply depends on B , resulting in the observed MR. Indeed, it is seen from Fig. 1(b) that for $\beta \gtrsim \beta_0$, $S_B \rightarrow 0$, resulting in sharp negative MR

$$\frac{\delta\rho_{xx}}{\rho} \sim \int_0^\infty \frac{dr}{l} \int_0^{a/r} d\phi e^{-2r/l} (e^{nS_B} - e^{nS_0}). \quad (3)$$

The following qualitative explanation of the observed linear MR was presented in Ref. [15]. The value $n(S_B - S_0)$ was estimated for $\phi = 0$ [Fig. 1(c)] to the first order in B as $-nr^3/R_c = -r^3/2alR_c$, where R_c is the cyclotron

radius. Assuming that this estimate also works at $\phi \neq 0$ and expanding $e^{nS_B} - e^{nS_0}$ to the first order in B , one gets $\delta\rho_{xx}/\rho \sim -l/R_c = -\omega_c\tau$.

In fact, the physical picture of the phenomenon is more subtle. The contribution of any trajectory with $\phi \neq 0$ is cancelled in the first order in B by the contribution of the time-reversed trajectory, since the values of $S_B - S_0$ are opposite for these paths [Fig. 1(d),(e)]. The cancellation does not occur only at very small $\phi \sim \beta$. The integration in Eq. 3 over $\phi < \beta$ yields $\delta\rho_{xx}/\rho \sim -\beta^2/\beta_0$. Larger values of ϕ also give a quadratic in β contribution to the MR. This contribution is positive and comes from the second order term in the expansion of $e^{nS_B} - e^{nS_0}$ in B . It follows from our results [Eqs. (1),(9)] that the contribution of small angles is dominant resulting in a negative parabolic MR [22]. We find that the parabolic MR crosses over to linear at very small $\beta \approx 0.05\beta_0$, which explains why the parabolic MR was not seen in numerical simulations [15] and experiment [16].

Next we sketch our calculations. We consider the Lorentz gas at $T = 0$, assuming that $\beta \ll 1$, $\beta_0 \ll 1$. In this case [23], $\delta\rho_{xx}/\rho = -(D - D_0)/D_0$, where $D_0 = v_F l_{tr}/2$ is the Drude diffusion coefficient for $B = 0$, $l_{tr} = 3l/4 = 3/8na$ is the momentum relaxation length and D is given by

$$D = \frac{1}{2} \int_0^\infty dt \langle \mathbf{v}(0) \mathbf{v}(t) \rangle = \frac{1}{2} \int d\mathbf{r} d\mathbf{v} \langle G \rangle \mathbf{v} \mathbf{v}_0. \quad (4)$$

Here $G = G(\mathbf{v}, \mathbf{v}_0, \mathbf{r})$ is the Fourier transform (at $\omega = 0$) of the retarded Green's function of the Liouville equation and $\langle \dots \rangle$ stands for the averaging over the positions of the disks. The equation for G reads

$$(\hat{L}_0 - \hat{T}_- - \hat{T}_+) G = \delta(\mathbf{r}) \delta(\mathbf{v} - \mathbf{v}_0), \quad (5)$$

where $\hat{L}_0 = \mathbf{v} \partial / \partial \mathbf{r} - \omega_c [\mathbf{v} \times \partial / \partial \mathbf{v}]$ is the Liouville operator of the free motion in the magnetic field. The interaction with disks is written in Eq. 5 in the form of a collision integral [18, 24]. The scattering operators $\hat{T}^\pm$ transform arbitrary function $f(\mathbf{r}, \mathbf{v})$ as follows,

$$\begin{aligned} \hat{T}^+ f(\mathbf{r}, \mathbf{v}) &= p_F \int d\mathbf{v}' \sigma(\varphi) \delta(\epsilon - \epsilon') n^+ f(\mathbf{r}, \mathbf{v}'), \\ \hat{T}^- f(\mathbf{r}, \mathbf{v}) &= -p_F \int d\mathbf{v}' \sigma(\varphi) \delta(\epsilon - \epsilon') n^- f(\mathbf{r}, \mathbf{v}), \end{aligned} \quad (6)$$

where $n^\pm = \sum_i \delta(\mathbf{r} - \mathbf{R}_i \pm \mathbf{a})$. Here $\mathbf{R}_i$ are the positions of the disks, delta-function $\delta(\epsilon - \epsilon')$ provides the energy conservation, p_F is the Fermi momentum, $\sigma(\varphi) = (a/2)|\sin(\varphi/2)|$ is the differential cross-section of one disk and φ is the angle between $\mathbf{v}'$ and $\mathbf{v}$. The vector $\mathbf{a} = \mathbf{a}(\mathbf{v}', \mathbf{v}) = a(\mathbf{v}' - \mathbf{v})/\sqrt{2(v^2 - \mathbf{v}'\mathbf{v})}$ is pointing from the center of a disk to the scattering point at the disk surface [Fig. 2]. Physically, operator $\hat{T}^+$ describes influx of particles with velocity $\mathbf{v}$ at the point $\mathbf{R}_i - \mathbf{a}$, while operator $\hat{T}^-$ describes the outflux of particles with




FIG. 2: Scattering of a particle on a hard disk.





FIG. 3: Diagrams, corresponding to the process shown in Fig. 1(a). Diagram (a) does not take into account effect of “empty corridor” and should be renormalized by (b).

velocity $\mathbf{v}$ at the point $\mathbf{R}_i + \mathbf{a}$. The Boltzmann equation is obtained from Eq. 5 by averaging the Liouville operator over the positions of the disks, yielding $\langle \hat{T}^+ \rangle f(\mathbf{r}, \mathbf{v}) = p_F n \int d\mathbf{v}' \sigma(\mathbf{v}, \mathbf{v}') \delta(\epsilon - \epsilon') f(\mathbf{r}, \mathbf{v}')$, $\langle \hat{T}^- \rangle = -1/\tau$. Here $n = \langle n^\pm \rangle$ is the concentration of the disks and $1/\tau = nv2a$ is the inverse full scattering time. Introducing now $\delta \hat{T}^\pm = \hat{T}^\pm - \langle \hat{T}^\pm \rangle$ and writing a formal solution of (5), $\hat{G} = (\delta + \hat{L}_0 - \hat{T}^- - \hat{T}^+)^{-1}$ as a series in $\delta \hat{T}^\pm$, we get

$$\langle \hat{G} \rangle = \hat{G}_0 + \sum_{\alpha, \gamma = \pm} \hat{G}_0 \langle \delta \hat{T}^\alpha \hat{G}_0 \delta \hat{T}^\gamma \rangle \hat{G}_0 + \dots, \quad (7)$$

where $\hat{G}_0 = (\delta + \hat{L}_0 + 1/\tau - \langle \hat{T}^+ \rangle)^{-1}$ is the Green's function of the Boltzmann operator (here $\delta \rightarrow 0$). Eq. 7 gives a regular way to calculate correlations, which are absent in the Boltzmann picture.

Consider first the case $B = 0$. Substituting the first term in the right hand side of Eq. 7 into Eq. 4, we get $D = D_0$. The second term in Eq. 7 describes the memory-effect due to diffusive returns. As discussed above, the main contribution comes from returns after a single scattering. This process is described by the diagram Fig. 3(a). The dashed line corresponds to the pairings $\langle \delta \hat{T}^\alpha \delta \hat{T}^\gamma \rangle$ ($\alpha, \gamma = \pm$), external wavy lines to the diffusion propagators $\hat{G}_0$. The internal line corresponds to the Boltzmann propagator truncated at one scattering $\hat{G}_- \langle \hat{T}^+ \rangle \hat{G}_-$, where $\hat{G}_- = (L_0 + 1/\tau)^{-1}$ is the ballistic propagator and $\langle \hat{T}^+ \rangle$ stands for one scattering event ($\hat{G}_-$ are shown by solid lines and $\langle \hat{T}^+ \rangle$ by the cross). This diagram yields $\delta \rho / \rho = -\delta D / D = (2\beta_0/3) \ln(1/2\beta_0)$ [17, 18, 19, 20, 21]. The terms of the N -th order in Eq. 7 contain N pairings (N dashed lines) and are typically small as β_0^N . However, there is a series of diagrams, shown in Fig. 3(b), whose contribution is of the order β_0 [19]. The internal dashed lines in this series only contain pairings $\langle \delta \hat{T}^- \delta \hat{T}^- \rangle$. Summing the diagrams Fig. 3(b) to-

FIG. 4: Backscattering process is parameterized by the angles φ_0, φ_f . The magnetic field changes the backscattering angle $\phi = \phi_0 + \phi_f + r/R_c$. The solid (dashed) line in (a) represents electron trajectory for $B = 0$ ($B \neq 0$). Different processes contributing to MR are shown in (b)-(e).

gether with Fig. 3(a), one gets an exact equation

$$\frac{\delta \rho}{\rho} = \frac{nl_{tr}}{4l} \text{Re} \int_a^\infty \frac{dr}{r} e^{-2r/l} \int_0^{2\pi} d\varphi_0 \int_0^{2\pi} d\varphi_f \sigma(\varphi_0) \sigma(\varphi_f) (1 - e^{i\varphi_0})(1 - e^{i\varphi_f}) e^{nS_0(r, \phi_0 + \phi_f)} = \frac{2\beta_0}{3} \ln \left(\frac{C}{2\beta_0} \right), \quad (8)$$

instead of qualitative estimate Eq. 2. Here φ_0, φ_f are the scattering angles [Fig. 4(a)], $\phi_0 \approx (a/r) \cos(\varphi_0/2)$, $\phi_f \approx (a/r) \cos(\varphi_f/2)$ and $C \approx 1.8$. Thus, addition of the series Fig. 3(b) to Fig. 3(a) leads to the following renormalization: $\ln(1/2\beta_0) \rightarrow \ln(C/2\beta_0)$. Physically, the series Fig. 3(b) accounts for the effect of the “empty corridor”. The N -th order term in this series corresponds to $N-1$ term in the Taylor expansion of the $\exp(nS_0)$ in Eq. 8. Four terms in the product $(1 - e^{i\varphi_0})(1 - e^{i\varphi_f}) = 1 - e^{i\varphi_0} - e^{i\varphi_f} + e^{i(\varphi_0 + \varphi_f)}$ correspond to four combinations of $(\pm, \pm)$ at the ends of external dashed lines in the diagrams shown in Fig. 3. They are connected with four different types of correlation at a given point $\mathbf{r}$. The diagram $(+, +)$ [Fig. 4(b)] corresponds to the process, where an electron has two real scatterings on a disk placed at point $\mathbf{r}$. The diagram $(-, -)$ [Fig. 4(c)] does not correspond to any real scattering at point $\mathbf{r}$. It just allows us to calculate correctly the probability for an electron to pass twice the region of the size a around point $\mathbf{r}$ without scattering. To interpret the diagram $(+, -)$, note that in the Boltzmann picture, which neglects correlations, the following process is allowed. An electron scatters on a disk and later on passes through the region occupied by this disk without a scattering [Fig. 4(d)] The diagrams $(+, -)$ correct the Boltzmann result by subtracting the contribution of such unphysical process. Analogous consideration is valid for diagram $(-, +)$ shown in Fig. 4[e].

For $B \neq 0$ the sum of diagrams shown in Fig. 3 can be expressed as an integral over angles φ_0, φ_f (scattering

angles for $B = 0$). The only difference from Eq. 8 is that one should replace $S_0 \rightarrow S_B$. For $\beta \ll 1$ the overlap area can be calculated as $S_B(r, \phi) = \int_0^r dx h(x)$, where $h(x) \approx (2a - |\phi x - x^2/R_c|) \theta(2a - |\phi x - x^2/R_c|)$, θ is the Heaviside step function and $\phi = \phi_0 + \phi_f + r/R_c$ [Fig. 4(a)]. The value of $\delta\rho_{xx}/\rho$ is obtained from Eq. 8 by replacing e^{nS_0} to $e^{nS_B} - e^{nS_0}$. Introducing dimensionless variables $T = r/l$, $z = \beta/\beta_0$ we get Eq. 1, where function $f(z)$ is given by

$$f(z) = \frac{3}{32} \int_0^\infty \frac{dT}{T} e^{-2T} \int_0^{2\pi} d\varphi_0 \int_0^{2\pi} d\varphi_f \cos\left(\frac{\varphi_0 + \varphi_f}{2}\right) \sin^2\left(\frac{\varphi_0}{2}\right) \sin^2\left(\frac{\varphi_f}{2}\right) (e^{sz} - e^{s_0}). \quad (9)$$

Here

$$s_z = \int_0^T dt \left(1 - \left|\zeta t - \frac{zt^2}{2}\right|\right) \theta\left(1 - \left|\zeta t - \frac{zt^2}{2}\right|\right), \quad \zeta = \frac{\cos(\varphi_0/2) + \cos(\varphi_f/2)}{2T} + \frac{zT}{2}, \quad s_0 = s_{z \rightarrow 0}. \quad (10)$$

Function $f(z)$ has the following asymptotics

$$f(z) = \begin{cases} 0.33z^2 & \text{for } z \lesssim 0.05 \\ 0.032(z - 0.04) & \text{for } 0.05 \lesssim z \lesssim 2 \\ 0.39 - 1.3/\sqrt{z} & \text{for } z \rightarrow \infty. \end{cases} \quad (11)$$

Note that there is a parametrically small nonanomalous correction to Eq. 1 due to returns after multiple scatterings, $\delta\rho'_{xx}/\rho \approx -0.2\beta_0\beta^2$ [15]. To compare the results of simulations [15] with the theoretical results in a wider region of parameters β, β_0 , we subtract $\delta\rho'_{xx}/\rho$ from the numerical curves. Theoretical and numerical [15] results are plotted in Fig. 5. in the universal units, $\delta\rho_{xx}/\rho\beta_0$ versus $z = \beta/\beta_0$. It is seen, that the theoretical and numerical results are in a very good agreement. The comparison with the experiment [16] is more difficult, because of the 50 % uncertainty in the sizes of the antidots. However, a good agreement with the experiment can be achieved by appropriate choice of a in the uncertainty interval [15].

Note finally that we fully neglected quantum effects. This is possible when $a \gg \sqrt{\lambda_F l}$ (λ_F is a Fermi wavelength). This criterion ensures that diffraction effects on the edges of the disks are not relevant at the scales of the order of l . In the opposite case, $a \ll \sqrt{\lambda_F l}$, the diffraction should destroy the “corridor effect”, does suppressing the anomalous MR. The detailed analysis of quantum effects will be presented elsewhere.

In summary, we have proposed a theory of the negative anomalous MR in the Lorenz gas. The analytical expression for the MR [Eqs. (1), (9), (11)] has been derived.

We thank M.I. Dyakonov for insightful discussions and R. Jullien for providing us with the numerical data. We are also grateful to I.V. Gornyi and D.G. Polyakov for useful comments. The work was partially supported by RFBR and INTAS.




FIG. 5: The value of $\delta\rho_{xx}/\rho\beta_0$ from Eqs. (1), (9) (solid line) shown as a function of β/β_0 together with the results of numerical simulations [15] presented for different values of β_0 (triangles for $\beta_0 = 0.09$, boxes for $\beta_0 = 0.06$, circles for $\beta_0 = 0.03$). Data for all numerical curves are shown for $\beta < 0.3$. Inset: The crossover from quadratic to a linear dependence at $\beta/\beta_0 \sim 0.05$. This crossover was not resolved in numerical simulations.

-
- [1] P.A. Lee and T.V. Ramakrishnan, Rev. Mod. Phys., **57**, 287 (1985).
 - [2] E.M. Baskin, L.N. Magarill, and M.V. Entin, Sov. Phys. JETP **48**, 365 (1978).
 - [3] D. Polyakov, Sov. Phys. JETP **63**, 317 (1986).
 - [4] A.V. Bobylev, F.A. Maaø, A. Hansen, and E.H. Hauge, Phys. Rev. Lett. **75**, 197 (1995).
 - [5] M. Fogler, A. Dobin, V. Perel, and B. Shklovskii, Phys. Rev. B **56**, 6823 (1997).
 - [6] M.M. Fogler and B.I. Shklovskii, Phys. Rev. Lett. **80**, 4749 (1998).
 - [7] A.V. Bobylev, F.A. Maaø, A. Hansen, and E.H. Hauge, J. Stat. Physics, **87**, 1205 (1997).
 - [8] E.M. Baskin and M.V. Entin, Physica B, **249**, 805 (1998).
 - [9] A. Kuzmany and H. Spohn, Phys. Rev. E, **57**, 5544 (1998).
 - [10] A.D. Mirlin, J. Wilke, F. Evers, D.G. Polyakov, and P. Wolffe, Phys. Rev. Lett. **83**, 2801 (1999).
 - [11] A.D. Mirlin, D.G. Polyakov, F. Evers, and P. Wolffe, Phys. Rev. Lett. **87**, 126805 (2001).
 - [12] D.G. Polyakov, F. Evers, A.D. Mirlin, and P. Wolffe, Phys. Rev. B **64**, 205306 (2001).
 - [13] D.G. Polyakov, F. Evers, and I.V. Gornyi Phys. Rev. B, **65**, 125326 (2002).
 - [14] A. Dmitriev, M. Dyakonov, and R. Jullien, Phys. Rev. B **64**, 233321 (2001).
 - [15] A. Dmitriev, M. Dyakonov, and R. Jullien, Phys. Rev. Lett. **89**, 266804 (2002).
 - [16] G. Gusev, P. Basmaji, Z. Kvon, L. Litvin, Yu. Nastaushev, and A. Toporov, Surface Science, **305**, 443 (1994).
 - [17] J. Dorfman and E. Cohen, Phys. Lett. **16**, 124 (1965).
 - [18] J. Van Leeuwen and A. Weiland, Physica, **36**, 457 (1967).
 - [19] J. Van Leeuwen and A. Weiland, Physica, **38**, 35 (1968).
 - [20] E.H. Hauge, in *Transport Phenomena*, edited by G. Kirzenow and J. Marro, Lecture Notes in Physics, No.31 (Springer, New York, 1974), p.337.
 - [21] R. Peierls, *Surprises in Theoretical Physics* (Princeton

- University, Princeton, 1979).
- [22] For very small β contribution of Lyapunov trajectories becomes important. This contribution is parametrically small compared to Eq.(1) for $\beta \gtrsim \beta_0^2$.
- [23] For $\beta \ll 1$ one can neglect the off-diagonal part of the conductivity tensor.
- [24] R. Zwanzig, Phys.Rev. B, **129**, 486, (1963).